

Semiclassical Electromagnetic Casimir Self-Energies

Martin Schaden*

Rutgers University, 211 Smith Hall, 101 Warren St., Newark, NJ 07102, U.S.A.

The electromagnetic Casimir energies of a spherical and a cylindrical cavity are analyzed semiclassically. The field theoretical self-stress of a spherical cavity with ideal metallic boundary conditions is reproduced to better than 1%. The subtractions in this case are unambiguous and the good agreement is shown that finite contributions from the exterior of the cavity are small. The semiclassical electromagnetic Casimir energy of a cylindrical cavity on the other hand vanishes to any order in the *real* reflection coefficients, whereas the field theoretic Casimir energy of a cylindrical boundary that is perfectly metallic and infinitesimally thin is finite and negative[18]. Contrary to the spherical case and in agreement with Barton's perturbative analysis[36], the subtractions in the spectral density for the cylinder are not universal when only the interior modes are taken into account[47]. The Casimir energy of a cylindrical cavity therefore depends sensitively on the physical nature of the boundary in the ultraviolet whereas the Casimir energy of a spherical one does not. The extension of the semiclassical approach to more realistic systems is sketched.

1. INTRODUCTION

Demonstrating that the collective interaction of atomic systems may have macroscopic consequences, Casimir obtained the now famous attractive force between two neutral metallic plates[1] from the boundary conditions they impose on the electromagnetic field. Half a century later, his prediction has been verified experimentally[2] to better than 1%.

Twenty years after Casimir's result for two parallel plates, Boyer calculated the zero-point energy of an ideal conducting spherical shell[3]. Contrary to intuition derived from the attraction between two parallel plates, the spherical shell tends to expand. Boyer's result has since been improved in accuracy and verified by a number of field theoretic methods[4,5,6,7,8] – even though there may be little hope of observing this effect experimentally in the near future[9].

Since field theoretic methods require explicit or implicit knowledge of cavity *frequencies*, they have predominantly been successfully employed to obtain the Casimir energies of classically *integrable* systems. Thus, in addition to a spherical cavity, the electromagnetic Casimir energies of dielectric slabs[10,11,12],

metallic parallelepipeds[13,14,15,16,17] and long cylinders[7,18,19,20,21,22,23] have been computed in this manner.

However, most systems are not integrable and often cannot even be approximated by such systems. It thus is desirable to develop reliable methods for estimating the Casimir energies of classically non-integrable and even chaotic systems. Balian, Bloch and Duplantier calculate Casimir energies based on a multiple scattering expansion to the Green's function[5,13,24]. This approach does not require knowledge of the quantum mechanical spectrum and the geometric expansion is exact for sufficiently smooth and ideally metallic cavities. Ultra-violet divergent contributions have to be subtracted at every order and the relative importance of the finite remainders is hard to assess a priori. It is in practice often difficult to carry the (in principle exact) expansion beyond the first few terms without semiclassical approximation². In[25] a semiclassical method based on Gutzwiller's trace formula[26] for the response function was proposed to estimate (finite) Casimir energies. It is appropriate

*email: mschaden@andromeda.rutgers.edu

²Restricting to just two reflections, the Casimir energy of an idealized spherical cavity of radius R is of the correct sign and quite accurately estimated[5] as $3\hbar c/(64R) \sim 0.0469\hbar c/R$

ate for Casimir energies of hyperbolic and chaotic systems[25,27,28] with isolated classical periodic orbits. The semiclassical approximation to the oscillating part of the spectral density of integrable systems was derived in[13,29] and has been used to estimate the force between concentric metallic cylindrical shells[30]. The agreement with exact results is rather impressive even when the radius of the outer cylinder is about four times that of the inner one[30]. Although not exact in general, the semiclassical approximation associates the finite (Casimir) part of the vacuum energy with optical properties of the system. It captures aspects of Casimir energies that have been puzzling for some time[31] and appears to be a reasonable approximation in the experimentally accessible regime where these forces are relatively large[25,31]. Path integral methods[32,33,34,35] in principle allow one to obtain Casimir interactions between disjoint bodies to arbitrary precision. Due to unresolved renormalization issues, these methods have so far not been used to study the self stress of cavities. The purpose of this article is to revisit and analyze the Casimir stress of (integrable) spherical and cylindrical cavities semiclassically and to compare it with field theoretic results.

The simplicity, transparency and somewhat surprising accuracy of the semiclassical approximation is demonstrated in Boyer's problem[3,4,5,6,7,8], that is in determining the electromagnetic Casimir energy of a spherical cavity with an (ideal) metallic boundary. Instead of directly proceeding from the semiclassical expressions for the oscillating part of the spectral density[13,29], our starting point will be the dual representation of the Casimir energy of integrable systems in terms of periodic orbits[26,29]. The subtractions in the spectral density that render the Casimir energy finite then are apparent and we can better analyze the semiclassical approximation. Even so, the semiclassical analysis of Boyer's problem is an order of magnitude simpler than any given previously. However, since no bounds are obtained, it at present is not possible to judge the accuracy of the result without comparing to field theoretic calculations[6]. It will become rather clear though, that the semiclassical analysis is accurate

enough to infer the sign of the Casimir energy of a cavity by geometric arguments if the contribution from periodic orbits does not vanish.

In ref.[30] periodic orbits in fact were found not to contribute to the Casimir energy of a long cylindrical cavity. The optical phases that give a positive semiclassical Casimir energy for the spherical cavity, lead to a vanishing one in the cylindrical case. The somewhat intriguing result that the Casimir energy of a cylinder[7,21,24,36,37,38] vanishes to first order in the reflection coefficients thus apparently is readily explained by geometric optics. However, the semiclassical Casimir energy of a cylindrical cavity vanishes to all orders in the *real* reflection coefficients and thus also vanishes for an ideal metallic cavity. The discrepancy to the finite field theoretic Casimir energy of an idealized, infinitesimally thin cylindrical boundary between non-dispersive media with the same speed of light[18,19,21] will be traced to the presence of a logarithmic divergence observed by Barton[36] in his perturbative treatment of the non-ideal dilute case. We will argue that the exact cancellation of this divergence in the field theoretic approach is due to the infinitesimal thickness of the boundary – interior and exterior contributions to the Casimir energy in this case depend on only one common scale – the radius of the cylinder.

2. The Dual Picture: Casimir Energies in Terms of Periodic Rays

Integrable systems may be semiclassically quantized in terms of periodic paths on invariant tori[39] – in much the same manner as Bohr first quantized the hydrogen atom. Although in general not an exact transformation, classical periodic orbits on the invariant tori are *dual* to the mode frequencies in the semiclassical sense. Applying Poisson's summation formula, the semiclassical Casimir energy (SCE) due to a massless scalar may be written in terms of classical periodic orbits[26,29,31],

$$\mathcal{E}_c = \frac{1}{2} \sum_{\mathbf{n}} \hbar \omega_{\mathbf{n}} - \text{UV subtractions}$$

$$\sim \frac{1}{2\hbar^d} \sum'_{\mathbf{m}} e^{-\frac{i\pi}{2}\beta_{\mathbf{m}}} \int_{sp} d\mathbf{I} H(\mathbf{I}) e^{2\pi i \mathbf{m} \cdot \mathbf{I} / \hbar} . \quad (1)$$

The components of the d -dimensional vector $\mathbf{I}$ in Eq.(1) are the actions of a set of properly normalized action-angle variables that describe the integrable system. The exponent of the integrand in Eq.(1) is the classical action (in units of $\hbar$) of a periodic orbit that winds m_i times about the i -th cycle of the invariant torus. $H(\mathbf{I})$ is the associated classical energy and $\beta_{\mathbf{m}}$ is the Keller-Maslov index[40,41] of a class of periodic orbits identified by the vector of integers $\mathbf{m}$. The latter is a topological quantity that does not depend on the actions $\mathbf{I}$. To leading semiclassical order, the (primed) sum extends only over those sectors $\mathbf{m}$ with classical periodic paths of finite action (see below). The correspondence in Eq.(1) can only be argued semiclassically[26,29] and the integrals on the RHS therefore should be evaluated in stationary phase approximation (*sp*).

Contributions to the Casimir energy from high frequencies correspond to those from short periodic orbits in this dual picture. Divergences due to periodic classical paths of vanishing length (and thus vanishing action) on the RHS of Eq.(1) are related to ultra-violet divergences of the mode sum on the LHS of Eq.(1). If these divergences can be subtracted unambiguously[31,36,42], the dependence of the vacuum energy on macroscopic properties of the system is semiclassically represented by contributions due to classical periodic orbits of finite action only. The primed sum on the RHS of Eq.(1) indicates this restriction³. The (divergent) Weyl contribution to the vacuum energy from the $\mathbf{m} = (0, \dots, 0)$ -sector in particular has to be subtracted. Together with an evaluation of the integrals in stationary phase, this defines the semiclassical Casimir energy (SCE). To physically interpret the SCE, one has to consider the implicit subtractions in the spectral density[24,31,36] in greater detail.

3. The Spherical Cavity

The semiclassical spectrum of a massless scalar is exact for a number of manifolds without boundary[43] and the definition of the SCE by the RHS of Eq.(1) coincides with the Casimir energy of zeta-function regularization in these cases. It also is exact for massless scalar fields satisfying periodic-, Neumann- or Dirichlet- boundary conditions on parallelepipeds[14,16,31] as well as for some tessellations of spheres[31,44,45]. In [25] the semiclassical approximation was argued to give the leading asymptotic behavior of the Casimir energy whenever the latter diverges as the ratio of two relevant lengths vanishes. All these criteria do not apply to the Casimir self-stress of a spherical cavity first considered by Boyer[3]. The latter is an integrable system, but the semiclassical spectrum is only asymptotically correct[13]. There furthermore is no ratio of lengths in which one might hope to obtain an asymptotic expansion. One therefore cannot expect the semiclassical approximation to be exact in this case. It nevertheless turns out to be quite accurate. The SCE is obtained by performing the integrals of Eq.(1) in stationary phase and has a very transparent interpretation in terms of periodic orbits *within* the cavity only. The sign of the SCE of a spherical cavity in particular will be quite trivially established and the good agreement supports the conjecture that the contribution from exterior modes mainly serves to cancel the ultra-violet divergence of the contribution from interior modes in the field theoretic approach[46,47]. The observed discrepancy of 1% compared to the field-theoretic result probably can be attributed to the error in the semiclassical estimate of low-lying eigenvalues of the Laplace operator – which is of similar size. Since boundary conditions for the electromagnetic field never are ideal, many corrections of similar and even greater magnitude would be required in any realistic situation.

The electromagnetic Casimir energy of any closed cavity with a smooth and perfectly metallic boundary may be decomposed into the contribution from two massless scalar fields – one satisfying Dirichlet's, the other satisfying Neumann's boundary condition on the surface[24].

³This is conceptually not so different from considering only the contribution of topologically non-trivial "instanton" sectors to the vacuum energy of a field theory.

Because the surface is ideally metallic, contributions to the spectral density from arbitrarily short closed paths that reflect off (either side of) the smooth surface cancel each other. Semiclassically there is no (potentially divergent) local contribution to the Casimir energy from such an idealized surface in the electromagnetic case – the local surface tension in fact vanishes for an infinitesimally thin surface[24,48]. Note that this cancellation is quite special for the electromagnetic field and metallic boundaries. It in general does not occur for a massless scalar field satisfying Neumann or Dirichlet boundary conditions on a spherical surface of arbitrary thickness or in even dimensions[46,49,50,51]. However, it was shown in [46] that the Casimir energy due to a scalar is finite for infinitesimally thin spherical cavities in odd dimensional spaces. The following argument illustrates the absence of ultraviolet divergent contributions proportional to the "area" of any *infinitesimally thin* even-dimensional surface if the speed of light in the surrounding medium is constant. Barring other scales and with a dimensionless cutoff⁴, the local contribution to the surface divergence from a small $((d-1)$ -dimensional) surface element dA is proportional to $\hbar c dA \sum_i f_i(\{R_j/R_k\})/R_i^d$, where R_i are (local) radii of curvature and the f_i are dimensionless functions of their ratios only. The surface divergence semiclassically arises due to arbitrary short closed optical paths on either side of the infinitesimally thin boundary that touch the surface at a single point only. But the radii of curvature on either side of a point on the infinitesimally thin surface are of equal magnitude and of opposite sign. This implies that divergent contributions from the two arbitrarily short closed optical paths with a common point on the surface that lie on either of its sides cancel precisely for odd d , whereas they are equal in even d – provided the boundary conditions and speed of light are the same for *both* sides of the surface. One therefore expects the scalar Casimir energy of such cavi-

ties to be analytic in the vicinity of odd d , but to in general diverge for even dimensional spaces. The finite Casimir energy of thin spherical cavities in d dimensions[46] as well as the perturbative analysis of such surface divergences[51] agree with this semiclassical analysis. In three dimensions, the Casimir energy due to a scalar field was found to be finite[19,47,52] also for an infinitesimally thin cylindrical cavity. The above argument is quite general and indicates that surface divergences in will cancel locally only for *infinitesimally thin* (and sufficiently smooth) even-dimensional boundaries when the speed of light is the same on either side. This cancellation of surface divergences is due to the *infinitesimal thinness* of the boundary and does not depend on the precise nature of the (frequency-independent) boundary condition. [It is quite different from the semiclassical cancellation of surface divergences of the electromagnetic field on ideal metallic boundaries in three dimensions. The latter occurs due to the different boundary conditions for the two scalar modes and does not depend on the thickness of the metallic surface.]

The only subtraction in the spectral density required for a finite Casimir energy in the electromagnetic case with idealized metallic boundary conditions thus is the Weyl contribution proportional to the volume of the sphere. The latter corresponds to ignoring the $\mathbf{m} = (0,0,0)$ contribution to the sum in Eq.(1). The remaining difficulty in calculating the SCE of an integrable system is a convenient choice of action-angle variables. For a massless scalar in three dimensions satisfying boundary conditions with spherical symmetry, an obvious set of actions is the magnitude of angular momentum, $I_2 = L$, one of the components of angular momentum $I_3 = L_z$ and an action I_1 associated with the radial degree of freedom.

Since the azimuthal angle of any classical orbit is constant, the energy $E = H(I_1, I_2)$ of a massless particle in a spherical cavity of radius R does not depend on $I_3 = L_z$. In terms of the previous choice of actions, the classical energy is implicitly

⁴In dimensional regularization this cutoff is just the (small) deviation from integer d and in the semiclassical context it is the ratio of the cutoff in the length of the shortest optical paths to some characteristic (fixed) length in the problem.

given by,

$$\pi I_1 + I_2 \arccos\left(\frac{cI_2}{ER}\right) = \frac{ER}{c} \sqrt{1 - \left(\frac{cI_2}{ER}\right)^2}. \quad (2)$$

The branches of the square root and inverse cosine in Eq.(2) are chosen so that I_1 is positive. It is convenient to introduce dimensionless variables

$$\lambda = 2ER/(\hbar c) \quad \text{and} \quad z = cI_2/(ER), \quad (3)$$

for the total energy (in units of $\hbar c/(2R)$) and the angular momentum (in units of ER/c) of an orbit. Note that $z \in [0, 1]$ and that the semi-classical regime formally corresponds to $\lambda \gg 1$, i.e. to wavelengths that are much shorter than the dimensions of the cavity. With the help of Eq.(2) and the definitions of Eq.(3), the semi-classical expression in Eq.(1) for the Casimir energy of a massless scalar field satisfying Neumann or Dirichlet boundary conditions on a spherical surface becomes,

$$\mathcal{E} = \frac{\hbar c}{4\pi R} \sum_{m,n \geq 0}' \Re \left[e^{-i\frac{\pi}{2}\beta(n,m)} \times \int_0^\infty d\lambda \lambda^3 \int_0^1 dz z \sqrt{1-z^2} e^{i\lambda[n(\sqrt{1-z^2}-z \arccos(z))+m\pi z]} \right]. \quad (4)$$

The integral over I_3 has here been performed in stationary phase approximation. Because the Hamiltonian does not depend on I_3 , only periodic orbits with $m_3 = 0$ contribute. Since $-I_2 \leq I_3 \leq I_2$, one has that $\int dI_3 = 2I_2 = \hbar \lambda z$. The factor $2I_2$ accounts for the $2(l+1/2)$ -degeneracy of states with angular momentum $L = \hbar(l+1/2) = I_2$. By taking (4 times) the real part in Eq.(4) one can restrict the summations to non-negative integers and choose the positive branch of the square root- and inverse cosine- functions in the exponent⁵. The Keller-Maslov index $\beta(n, m)$ of a classical sector depends on whether Neumann or Dirichlet boundary conditions are satisfied on the spherical shell and will shortly be determined.

For positive integers m and n , the phase of the integrand in Eq.(4) is stationary at $z = \bar{z}(n, m) \in$

⁵The primed sum now implies half the summand if one of the integers vanishes as well as the absence of the $m = n = 0$ term.

$[0, 1]$ where,

$$\begin{aligned} 0 &= -n \arccos(\bar{z}) + m\pi \\ \Rightarrow \bar{z}(n, m) &= \cos(m\pi/n), \quad n \geq 2m > 1. \end{aligned} \quad (5)$$

Restrictions on the values of m and n arise because $\arccos(\bar{z}) \in [0, \pi/2]$ on the chosen branch. The phase is stationary at classically allowed points only for sectors with $n \geq 2m > 1$. Semi-classical contributions to the integrals of other sectors arise due to the endpoints of the z -integration at $z = 0$ and $z = 1$ only. Such "diffractive" contributions are of sub-leading order in the asymptotic expansion of the spectral density for large λ . Note that $m \rightarrow m+n$ amounts to the choice of another branch of the inverse cosine.

The classical action in sectors with stationary points is,

$$\begin{aligned} S_{cl}(n, m) &= \hbar \lambda n \sin(m\pi/n) \\ &= (E/c) 2nR \sin(m\pi/n) = (E/c) L(n, m), \end{aligned} \quad (6)$$

where $L(n, m)$ is the total length of the classical orbit. Some of these periodic orbits are shown in Fig. 1. The integer m in Eq.(6) gives the number of times an orbit circles the origin. The integer $n > 1$ gives the number of times an orbit touches the spherical surface. As indicated in Fig. 1, the set of classical periodic orbits in the (n, m) -sector form a caustic surface and a double covering is required for a unique phase-space description[40]. The two sheets are joined at the inner caustic [indicated by a dashed circle in Fig. 1] and at the outer spherical shell of radius R . Every orbit that passes the spherical shell n times also passes through the caustic n times. The cross-section of a bundle of rays is reduced to a point at the spherical caustic surface. The caustic thus is of second order and is associated with a phase loss of π every time it is crossed. At each specular reflection off the outer shell, Dirichlet boundary conditions require an additional phase loss of π whereas there is no change in phase for Neumann boundary conditions. Altogether the Keller-Maslov index of sector (n, m) depends on n only and is given by,

$$\beta(n, m) = \begin{cases} 0, & \text{for Dirichlet b.c.} \\ 2n, & \text{for Neumann b.c.} \end{cases}. \quad (7)$$

Since the electromagnetic Casimir energy of a cavity with a smooth surface on which (ideal) metallic boundary conditions hold can be viewed as due to two massless scalar fields, one satisfying Dirichlet and the other Neumann boundary conditions[24] on this surface, only sectors (n, m) with *even* $n = 2k \geq 2m \geq 2$ contribute[24] to the SCE in leading order of the asymptotic expansion for large λ .




Fig.1: Classical periodic rays within a ball or a solid cylinder. a) The shortest primitive rays with winding numbers $(n, m) \in \{(2, 1), (3, 1), (4, 1)\}$. b) Primitive rays to winding numbers $(n, m) = (5, 1)$ and $(5, 2)$. Caustic surfaces are shown as thin circles. The dashed part of any trajectory is on one sheet and its solid part on the other of a two-sheeted covering space. The "phase space" of the $(5, 2)$ sector is indicated by the hatched area. Note that caustics are of 2nd order for a spherical cavity but of 1st order for a cylindrical one.

Note that sectors with $m = 0$ or $n = 0$ have vanishing classical action and do not contribute to the SCE. Eq.(5) implies that extremal paths in the $(n > 0, m = 0)$ sectors have maximal angular momentum $1 = \bar{z} = lc/(ER)$. These are great circles that are wholly within the spherical shell in a plane perpendicular to the one under consideration, i.e. classical orbits with $I_3 = L_z = 0$. Because the measure of the z -integral vanishes at $z = 1$ like $\sqrt{1-z}$ these classical paths are extremal but not stationary. This can also be seen by expanding the exponent in Eq.(4) about the stationary point $\bar{z}(n, m)$. For $(n > 0, m > 0)$ the second derivative of the classical action is finite

at $\bar{z}(n, m)$,

$$\frac{\partial^2}{\partial \bar{z}^2} [n(\sqrt{1-\bar{z}^2} - \bar{z} \arccos(\bar{z})) + m\pi \bar{z}] = \frac{n}{\sin(\frac{m\pi}{n})}, \quad (8)$$

whereas it diverges in sectors with $m = 0$. The behavior of the exponent for $z \sim 1$ in this case is,

$$\sqrt{1-z^2} - z \arccos(z) = \frac{2\sqrt{2}}{3}(1-z)^{3/2} + \mathcal{O}((1-z)^{5/2}). \quad (9)$$

Quadratic fluctuations about the classical orbit with $m = 0$ thus have vanishing width and these sectors do not contribute in stationary phase approximation. To leading semiclassical accuracy, the Casimir energy of a spherical cavity with an ideal metallic boundary therefore is,

$$\begin{aligned} \mathcal{E}_{\text{EM}}^{\text{ball}} &\sim \frac{\hbar c}{4\pi R} \text{Re} \sum_{n=1}^{\infty} (1^n + (-1)^n) \sum_{m=1}^{n/2} \\ &\times \int_0^{\infty} d\lambda \lambda^3 e^{in\lambda \sin(\frac{m\pi}{n})} \int_0^1 dz z \sqrt{1-z^2} e^{i \frac{n\lambda(z - \bar{z}(n,m))^2}{2 \sin(\frac{m\pi}{n})}} \\ &\sim \frac{\hbar c}{R} \left[\sum_{k=1}^{\infty} \frac{1}{16\pi k^4} + \sum_{k=2}^{\infty} \frac{15\sqrt{2}}{256k^4} \sum_{m=1}^{k-1} \frac{\cos(\frac{m\pi}{2k})}{\sin^2(\frac{m\pi}{2k})} \right] \\ &\sim 0.04668... \frac{\hbar c}{R}. \end{aligned} \quad (10)$$

This semiclassical estimate is only about 1% larger than the best numerical value[6] $0.04617... \hbar c/R$ for the electromagnetic Casimir energy of a spherical cavity with an infinitesimally thin metallic surface. The error is of the same order as that of limiting the multiple reflection expansion to just two reflections, which gives[5] $\mathcal{E}_{\text{EM}}^{\text{ball}} \sim 0.0469 \hbar c/R$. Note that the contribution from the $(2k, k)$ sectors had to be considered separately in Eq.(10) since the measure $dz z$ vanishes at the stationary point $\bar{z}(2k, k) = \cos(\pi/2) = 0$ of the integrand, which is an endpoint of the integration domain. As can be seen in Fig. 1a), the classical rays of $(2k, k)$ -sectors go back and forth between antipodes of the cavity and pass through its center – they have angular momentum $L = 0 = \bar{z}(2k, k)$.

The shortest primitive orbits give somewhat less than half $(1/(16\pi) \sim 0.02)$ of the total SCE of the spherical cavity – much less than the 92% they contribute to the Casimir energy of parallel plates. The main reason is that contributions

only drop off as $1/k^2$ rather than like $1/k^4$ as for parallel plates. The length of an orbit in the $(4, 1)$ -sector (the inscribed square in Fig. 1a) furthermore is just a factor of $\sqrt{2}$ longer than a $(2, 1)$ -orbit [which in turn is a factor of $1/\sqrt{2}$ shorter than a $(4, 2)$ -orbit]. To estimate the magnitude of the contribution from any particular sector one has to take the available phase space as well as the ray's length into account. Thus, although the length of a $(2k, 1)$ -orbit tends to $2\pi R$ for $k \rightarrow \infty$, the associated phase-space (essentially given by the volume of the shell between the boundary of the cavity and the inner caustic) decreases like $1/k^2$. This accounts for the relatively slow convergence of the sum in Eq.(10). To achieve an accuracy of 10^{-5} , the first 50 terms of the sum were evaluated explicitly and the remaining contribution was estimated using Richardson's extrapolation method.

4. The Cylindrical Cavity

The example of a spherical cavity shows that the SCE in some instances is surprisingly accurate. However, there evidently are systems without periodic classical orbits, such as the two perpendicular planes investigated in [34], or the Casimir pendulum of [53]. None of these systems is integrable, and although there are no stationary periodic classical rays, periodic rays of *extremal* (shortest) length do exist. Semiclassically, such extremal periodic rays are associated with *diffraction* [54,55]. The inclusion of diffractive contributions in the semiclassical estimate of Casimir energies has so far only been attempted for a system of spheres [56]. It will become evident below that diffractive contributions also play a central role in the Casimir energy of a cylindrical cavity.

The Casimir energy of a dilute cylindrical gas of atoms was found to vanish in [57]. A number of calculations have confirmed that there is no contribution up to second order in the reflection coefficients for dielectrics [36,37,38] and for media where the speed of light on either side of an infinitesimally thin cylindrical boundary is the same [7,20,21,22,24]. Balian and Duplantier even conjectured that the Casimir energy of an ideal metallic cylindrical cavity may vanish [24] to all

orders of the multiple reflection expansion. The non-vanishing Casimir energy of an ideal metallic cylindrical cavity [18] was reanalyzed in the framework of zeta-function regularization. It was confirmed that the Casimir energy of an ideal metallic cylinder only vanishes to leading order and that higher orders in the reflection coefficients all give a non-vanishing contribution [19]. However, some mathematical prowess is required to analytically prove the lowest order cancellation in the field-theoretic approach [37,38]. That a number of separate contributions should conspire to a null result without apparent physical reason has been considered by many as somewhat "mysterious" [7,12]. The suspicion that this cancellation could be a purely geometrical effect, is nourished by the fact that the finite part of the pair-wise Van Der Waals interaction energy of a dilute gas of atoms vanishes for a cylinder [36,57] but not for other geometries. In [30] it was found that the semiclassical contribution due to periodic rays also vanishes for a cylinder. However, a careful perturbative analysis reveals that the interaction energy of any real dilute cylindrical gas of atoms includes a logarithmic divergence in addition to divergent contributions proportional to the volume and surface area of the cylinder [36]. The subtraction of this logarithmic divergence generally is ambiguous and the Casimir energy of a cylindrical cavity depends sensitively on properties of its boundary [58] in the ultraviolet. A particular boundary (say an infinitesimally thin cylindrical shell separating media with the same speed of light) thus may have a finite (negative) Casimir energy, whereas a very small modification of this boundary (say finite thickness) leads to a logarithmic divergence.

The calculation below supports this possibility. It is already known from [30] that the semiclassical contribution to the Casimir energy vanishes for a cylinder. The SCE of a cylindrical cavity in fact vanishes to *all orders* in the reflection coefficients for the same reason that the SCE of a spherical cavity is positive – due to relatively obvious optical phases. The semiclassical point of view thus gives a straightforward and physically acceptable explanation for otherwise mysterious cancellations. It also indicates that any addi-

tional phase change at the boundary will destroy this delicate mechanism. The finite electromagnetic Casimir energy[18] of a cylinder with idealized metallic boundary conditions on the other hand is more difficult to explain semiclassically. However, contrary to a spherical cavity and in agreement with the perturbative result of [36], the semiclassical expression for the Casimir energy of a cylindrical cavity also is logarithmic divergent. This is due to "diffractive" end-point contributions that are ignored in stationary phase approximation. There is reason to believe[47,52] that the subtraction of this logarithmic divergence by the contribution from "exterior" modes gives a finite Casimir energy for an ideal metallic and infinitesimally thin cylinder[18].

To exhibit these effects we revisit the calculation of the electromagnetic SCE of a long cylindrical cavity, or rather of a very thin torus with one perimeter L that is much larger than the other, $L \gg 2\pi R$. This is an integrable system. In the limit $R/L \rightarrow 0$, the only classical trajectories of relevance are again those of Fig. 1 and the SCE of a long cylindrical cavity can be obtained along similar lines as that of a spherical one – with some important modifications. Due to the toroidal symmetry of the (long) cylinder, the third action $I_3 = Lp_L/(2\pi)$ in this case is proportional to the conserved momentum p_L along the axis of the (thin) cylinder and in Eq.(2) the energy E must be replaced by $\sqrt{E^2 - (2\pi cI_3/L)^2}$. The second action furthermore is the angular momentum rather than just its magnitude. It again is convenient to consider dimensionless quantities for the fraction $-1 \leq x \leq 1$ of the total momentum along the axis of the cylinder, for the ratio $-1 \leq z \leq 1$ of the angular momentum to the maximal possible angular momentum of a photon within the cavity and for its energy $0 \leq \lambda < \infty$ in units of $\hbar c/(2R)$,

$$\lambda = 2ER/(\hbar c), \quad z = \frac{cI_2/(ER)}{\sqrt{1-x^2}}, \quad x = \frac{2\pi cI_3}{EL}. \quad (11)$$

Proceeding as in the spherical case, the semiclassical expression in Eq.(1) for the SCE of a massless scalar field satisfying Neumann or Dirichlet boundary conditions on a cylindrical surface be-

comes,

$$\mathcal{E}_{\text{cyl}} = \frac{\hbar c L}{16\pi^2 R^2} \sum_{m,n \geq 0}' \Re \left[e^{-i\frac{\pi}{2}\beta(n,m)} \int_0^\infty d\lambda \int_{-1}^1 dz \int_{-1}^1 dx \times \right. \\ \left. \times \lambda^3 \sqrt{1-z^2} e^{i\lambda \sqrt{1-x^2} [n(\sqrt{1-z^2} - z \arccos(z)) + m\pi z]} \right]. \quad (12)$$

The contribution from periodic orbits that wind around the perimeter of the torus is negligible in the $R/L \rightarrow 0$ limit and has been omitted in Eq.(12). The phase of the integrand in Eq.(12) is stationary at $\bar{x} = 0$ (corresponding to $p_L = 0$) and $\bar{z}(n, m)$ given in Eq.(5). Since the domain of integration for the z -variable differs from the spherical case, sectors with $1 < m < n - 1$ have non-trivial stationary points. The classical action of an (n, m) -sector is the same as for the spherical cavity and is given by Eq.(6). The fluctuations about such a classical ray on the other hand are quite different for cylindrical and spherical cavities. To quadratic order in the fluctuations about the stationary point $\bar{x} = 0, \bar{z}(n, m)$, the action for the cylinder is

$$S_{nm} \sim n \left[\left(1 - \frac{x^2}{2}\right) \sin \frac{m\pi}{n} + \frac{(z - \bar{z}(n, m))^2}{2 \sin \frac{m\pi}{n}} \right]. \quad (13)$$

The unconstrained Gaussian integrals over $z - \bar{z}(n, m)$ and x result in a factor of $2\pi/(n\lambda)$ in stationary phase approximation. Note that the phases of $\pm\pi/4$ associated with the two Gaussian integrals cancel in this case. Performing also the integral over λ in Eq.(12) finally gives,

$$\mathcal{E}_{\text{cyl}} = \frac{\hbar c L}{4\pi R^2} \sum_{n=2}^\infty \sum_{m=1}^{n-1} \Re \frac{-i e^{-i\frac{\pi}{2}\beta(n,m)}}{n^4 \sin^2 \frac{m\pi}{n}}. \quad (14)$$

The crucial difference to a spherical cavity is the phase factor of $-i$. It arises because the fluctuations of a cylindrical system have one fewer zero-mode than those of a spherical one⁶. The

⁶The Hamiltonian of a spherical cavity does not depend on $I_3 \propto L_z$, whereas it does depend on $I_3 \propto p_L$ for the cylindrical cavity. The 2×2 Hessian matrix $H_{ij} = \partial^2 H / \partial I_i \partial I_j$ with $3 > i, j > 1$ has one zero mode for a spherical cavity, but none for the cylindrical geometry. This difference in zero modes implies[31] an additional phase loss of $\pi/2$ for the periodic rays of a cylindrical cavity.

additional phase loss of $\pi/2$ ultimately is responsible for the vanishing of the SCE of a cylindrical cavity. To verify this we only need to compute the Keller-Maslov index $\beta(n, m)$ for Neumann and Dirichlet boundary conditions. The caustics of the cylindrical cavity are of first order rather than second: the cross-section of a bundle of rays becomes one-dimensional at the caustic – it is focussed to a line rather than a point. Taking into account the phase retardation by $\pi/2$ every time a ray passes a first order caustic, the analogous result to Eq.(7) for a cylindrical cavity is,

$$\beta(n, m) = \begin{cases} 3n, & \text{for Dirichlet b.c.} \\ n, & \text{for Neumann b.c.} \end{cases} \quad (15)$$

Contributions from paths with Neumann and Dirichlet boundary conditions and an odd number of reflections cancel each other and, as for the spherical cavity, only sectors to even $n = 2k = 2, 4, \dots$ contribute to the electromagnetic SCE [for smooth metallic cavities this is quite generally so[24]]. Summing contributions to the electromagnetic Casimir energy from the two scalars in Eq.(14) then gives the null result

$$\mathcal{E}_{\text{cyl}}^{\text{EM}} = \frac{\hbar c L}{32\pi R^2} \sum_{k=1}^{\infty} \sum_{m=1}^{2k-1} \Re \frac{-i(-1)^k}{k^4 \sin^2 \frac{m\pi}{2k}} = 0. \quad (16)$$

In Eq.(16) *every* periodic orbit gives a vanishing contribution to the SCE of a cylindrical cavity. The cancellation evidently depends on a delicate relation between the optical phases. It is interesting that a small additional phase *loss* at each reflection off the surface results in a *negative* SCE for a cylindrical cavity, but that the Casimir energy vanishes as long as the above phase relations hold – even if the magnitude of the reflection coefficients is less than unity. The SCE in this sense is in line with previous results[36,37,38,57] for the Casimir energy of a dilute dielectric cylinder, and in fact supports the conjecture of Balian and Duplantier in[24]. The non-vanishing Casimir energy of a cylindrical cavity with an infinitesimally thin ideal metallic boundary on the other hand is not so easily explained by this semiclassical point of view.

Some insight is gained by noting that the contribution of *any* sector to the SCE of a cylin-

dricl cavity in Eq.(12) – even sectors with non-trivial periodic classical paths – diverges. This is in marked contrast to the spherical case, where the contribution from sectors with non-trivial periodic classical paths (characterized by $n \geq 2m > 1$) is *finite*. The divergence is readily made explicit by scaling $\lambda\sqrt{1-x^2} \rightarrow \lambda$ in the integral of Eq.(12). Without ultraviolet cutoff, the resulting x -integral in this case formally gives the factor,

$$\int_{-1}^1 \frac{dx}{(1-x^2)^2} \sim \infty, \quad (17)$$

which diverges due to the behavior of the integrand near the endpoints of the integral at $x = \pm 1$. It may be regulated by introducing an *ultraviolet* cutoff Ω of some sort for the energy integral [that is in the integral over λ]. As may be seen from Eq.(17), the regulated integral will always include terms that are *logarithmically* divergent as $\Omega \rightarrow \infty$. The subtraction of a logarithmic divergence depends on details of the cutoff and thus is sensitive to ultraviolet properties of the boundary[42]. The evaluation of (in principle divergent) integrals in stationary phase could be considered *one* way of subtracting this divergence. Because the divergence is logarithmic, the subtraction is by no means unique in this case. The presence of a logarithmic divergence for cylindrical cavities was first emphasized by Barton[36] in his perturbative treatment of a dilute gas of atoms. It also is evident in the contribution from interior modes to the Casimir energy of an ideally metallic cylinder[47].

The foregoing is compatible with previous results[18,21,7] that the Casimir energy of a cylindrical cavity is finite if the speed of light inside and outside its *infinitesimally thin* boundary are the same. It for instance is negative for idealized metallic boundary conditions[18]. The Casimir energy in this case apparently does not suffer from any logarithmic divergences (or equivalently, from any pole ambiguities in zeta function regularization). The Casimir energy is finite for the infinitesimally thin boundary, because the logarithmic divergent contribution from interior modes is precisely cancelled by the similarly logarithmic divergent contribution from exterior modes.

Since the boundary is infinitesimally thin and the speed of light on both sides of the boundary is the the same, a precise cancellation is possible. The divergence reappears for a dielectric cavity in vacuum with a lower speed of light in the dielectric[58]. This occurs for a spherical as well as for a cylindrical cavity, but with an important difference: the divergence in the spherical case is not logarithmic and may be unambiguously subtracted[9]. The subtraction of the logarithmic divergence in the Casimir energy of the cylindrical cavity on the other hand requires the introduction of some energy scale that describes ultraviolet properties of the boundary. An analogous problem would be encountered for an ideal metallic boundary of finite thickness[47] and in fact for almost any small deviation from an idealized and infinitesimally thin cylindrical boundary between two media with identical speed of light. Paradoxically, *defining* the Casimir energy of an cylindrical cavity in a manner that does not depend on detailed ultraviolet properties of its boundary appears all but impossible.

It perhaps is worth mentioning in this regard that the Casimir energy of a massless scalar excitation *on* the two-dimensional spherical or toroidal boundaries is well-defined. For a spherical shell and a very thin torus, this Casimir energy has the same dependence on the dimensions as the Casimir energies of the corresponding cavities. For a two-sphere (S_2) and a very thin torus T_2 with $L \gg 2\pi R$ these Casimir energies are

$$\begin{aligned} \mathcal{E}_{S_2} &= 0 \\ \mathcal{E}_{T_2} &= -\frac{\hbar c L}{4\pi^3 R^2} \zeta(3) \sim -0.0097 \dots \frac{\hbar c L}{R^2} . \end{aligned} \quad (18)$$

Note that these Casimir energies of a massless scalar on two-dimensional spherical and toroidal surfaces are exactly reproduced semiclassically[31,43]. The presence of scalar surface modes therefore does not change the Casimir energy of a spherical cavity but could very well contribute to that of a cylindrical one. The Casimir energy of a massless degree of freedom on a torus not only is of the same form, but also of the same sign and order of magnitude as the Casimir energy of an ideal metallic cylindrical cavity[18,19]. Such a contribution from

massless surface modes thus might be important for a cylindrical cavity and would be difficult to separate from the contribution due to cavity modes.

5. Discussion

The semiclassical approximation to the Casimir energy of a cavity to leading order includes only contributions from quadratic fluctuations about stationary periodic classical rays. Since all periodic rays lie in the interior, the SCE of a concave cavity to leading order depends on the exterior only indirectly through reflection coefficients. Periodic classical rays furthermore are of finite length. Their contribution to the Casimir energy thus is ultraviolet finite. However, this approximation is sensible only if UV-divergent contributions to the vacuum energy can be subtracted unambiguously from the spectral density. Logarithmically divergent contributions to the vacuum energy require a subtraction scale[42]. The latter is a clear indication that the subtraction cannot be universal since it depends sensitively on the UV-properties of the boundary. Small changes in the boundary conditions in this case do not necessarily correspond to small changes in the Casimir energy. The *local* properties of a boundary apparently include its thickness: whereas the Casimir energy of a cylindrical cavity with an ideal and infinitesimally thin metallic boundary is finite[18] to any order in the (real) reflection coefficients[19], a logarithmic dependence on the cutoff appears in more realistic situations[36,47]. An ambiguous subtraction also is required in the semiclassical approximation. The absence of any logarithmic divergence for the infinitesimally thin boundary apparently is due to a cancellation by exterior modes. Such a cancellation of logarithmic singularities can occur when exterior and interior modes depend on precisely the same scale, the radius R of the cylindrical cavity in this case. Although the two logarithmic divergences (each proportional to $\hbar c L/R^2$ for dimensional reasons) cancel in the idealized situation, they would not do so for a boundary of finite thickness.

We considered only the semiclassical Casimir energy (SCE) of a spherical and of a toroidal cav-

ity with ideal metallic boundary conditions, that is, with real reflection coefficients of unit magnitude. These are integrable systems and the SCE was obtained from the "dual" description of the spectral density in terms of periodic paths on invariant tori[26,29]. The winding numbers of a periodic orbit are dual to the quantum numbers of a mode. In stationary phase approximation the SCE of a spherical cavity is *positive* and reproduces the field theoretic value for an infinitesimally thin metallic boundary to within 1%. The calculation is rather short and straightforward and leads to the convergent sum of Eq.(10). Each term in this sum may be interpreted as the contribution from a class of periodic rays. A few of the shorter primitive periodic rays are depicted in Fig. 1. The contribution from any sector is finite in this case.

The contribution from periodic orbits to the SCE of a cylindrical cavity with an ideal metallic boundary on the other hand vanishes to all orders in the number of reflections[30]. This occurs due to an overall phase change by an odd multiple of $\pi/2$ for any classical periodic ray. Restricting to just two reflections, this null result agrees with field theoretic calculations for infinitesimally thin metallic boundaries[7,21,23,24,59]. The vanishing SCE appears to support the conjecture of Balian and Duplantier that the Casimir energy of a metallic cylindrical cavity may vanish. However, contrary to the spherical case, the contributions of *any* classical sector to the SCE of a cylindrical cavity diverges. Without subtraction of the UV-divergent part, the (finite) semi-classical contribution to the vacuum energy we obtained is not very meaningful. Unfortunately the divergence of the integral in Eq.(17) includes a logarithmic dependence on the cutoff. The subtraction of UV-divergent contributions to the Casimir energy of a cylinder thus is sensitive to a scale and cannot be achieved in a universal fashion. The logarithmic dependence on the cutoff was first observed by Barton[36] in his perturbative calculation of the vacuum energy for a (dilute) gas of cylindrical shape to lowest order in the fine structure constant. Semiclassically this would also correspond to considering the contribution from rays with only two reflections ($n = 2$). That

the UV-subtractions are fragile and depend crucially on the UV-properties of the boundary is also observed when the speed of light within and outside an infinitesimally thin cylindrical boundary differ[58]. In the electromagnetic case, the logarithmic divergences of exterior and interior contributions to the vacuum energy of a metallic cylinder cancel for an infinitesimally thin metallic boundary[47]. However, they in general cannot be unambiguously subtracted[52].

These examples of a spherical- and cylindrical cavity show that the SCE is quite reasonable and is rather simple to calculate when the Casimir energy is robust, that is, when the necessary subtractions depend on global properties of the system only and do not require the introduction of an additional scale. This apparently is not possible if the divergence is logarithmic[42,31]. The classical periodic paths that contribute to the SCE of a concave cavity in stationary phase approximation lie entirely within the cavity. Their contribution depends on the exterior of the cavity through reflection coefficients only. It has been argued for some time that a Casimir energy obtained without explicit inclusion of exterior modes (as for a parallelepiped[14,16,17]) is all but meaningless[12]. The criterion favored here[36,31] considers any definition of a Casimir energy reasonable (and in principle physically realizable) in which the UV-divergences of the vacuum energy have been subtracted without reference to local properties of the boundary. The subtraction may (and in general will) include divergent contributions from exterior modes. The Casimir energy of a parallelepiped can be considered a case in point: as Power[60] did for just two slabs, one can always assemble parallelepipeds to a cube with fixed dimensions – the Casimir energy of an individual parallelepiped[14,16,17] in this case reflects changes in the vacuum energy of the whole cube as the three dividing planes are moved adiabatically. By moving interior surfaces of the cube (that in principle could have finite thickness), one measures only the finite part of its vacuum energy that depends on the dimensions of the individual parallelepipeds. By contrast, it is difficult to imagine that global changes in a vacuum energy are measurable (or even physi-

cally relevant) if their finiteness depends critically on local characteristics of the system[31]. Perhaps somewhat surprisingly, the electromagnetic Casimir energy due to the interior of a very long cylindrical cavity does not appear to be robust in this sense, whereas the electromagnetic Casimir energy due to the interior of a spherical cavity is.

Apart from relating Casimir energies to optical properties, one of the advantages of a semiclassical description would be the possibility to model more realistic (but robust) physical systems. The previous considerations are readily extended to dielectrics by using appropriate complex and in general frequency-dependent reflection coefficients. In the case of dielectric slabs Milton has shown[12] that Lifshitz's theory[10,11] may be reproduced in this manner. Finite temperature is incorporated[61] by allowing periodic rays to also wrap around a fictitious periodic extra dimension of circumference $\hbar c/(kT)$. Finite temperature corrections thus are small if some classical periodic paths are much shorter than this circumference. At room temperature the length of a periodic ray increases by about 7.6 microns every time it winds about the temperature direction. Temperature corrections therefore are tiny for most nanometer scale experiments⁷ but could be of greater interest in some astrophysical considerations ($3^\circ K \equiv 1\text{mm}$). Corrections due to surface roughness generally will be more important in technological applications. Many classical models for diffuse reflection from rough surfaces exist and Lambert's Law is easily incorporated in the semiclassical approach by appropriate reflection coefficients. The dependence on the wavelength perhaps can be modelled by a term of the action that accounts for stochastic fluctuations in the length of a classical periodic orbit upon reflection from rough surfaces. Apart from an average change in length, this leads to a damping term of the form $-(\Delta L E/\hbar c)^2/2$ in the classical action, where $(\Delta L)^2$ is the variance in the length of the periodic orbit. Assuming that this variance is itself proportional to the length of the orbit, surface roughness can semiclassically perhaps be

⁷The claim that this correction has been measured to sufficient accuracy[2] to distinguish between different approaches has recently been disputed[62].

modelled by the modified dispersion

$$cp(E) = E + i\varepsilon E^2/(\hbar c), \quad (19)$$

where ε is a typical length scale for the (stochastic) roughness of the surface. The predominant effect of the modified dispersion of Eq.(19) is that contributions to the Casimir energy from wave lengths $\lambda \ll \varepsilon$ are very much suppressed. A similar conclusion may be drawn from a recent and considerably more sophisticated analysis[63].

Acknowledgements: I am indebted to Larry Spruch for advice and numerous discussions and would like to thank K. A. Milton for a critical reading of the manuscript. This work was partially supported by the National Science Foundation under Grant No. PHY0555580.

REFERENCES

1. H. B. G. Casimir, Kon. Ned. Akad. Wetensch. Proc. **51**, 793 (1948).
2. S. K. Lamoreaux, Phys. Rev. Lett. **78**, 5 (1997); U. Mohideen and A. Roy, Phys. Rev. Lett. **81**, 4549 (1998); T. Ederth, Phys. Rev. **A 62**, 062104 (2000); G. Bressi, G. Carugno, R. Onofrio and G. Ruoso, Phys. Rev. Lett. **88**, 041804 (2002); R. S. Decca, D. Lopez, E. Fishbach, G. L. Klimchitskaya, D. E. Krause, and V. M. Mostepanenko, Ann. Phys. **318**, 374 (2005).
3. T. H. Boyer, Phys. Rev. **174**, 1764 (1968).
4. B. Davies, J. Math. Phys. **13**, 1324 (1972).
5. Annals Phys. **104**, 300 (1977); *ibid* **112**, 165 (1978).
6. K. A. Milton, L. L. DeRaad and J. S. Schwinger, Annals Phys. **115**, 388 (1978).
7. G. Lambiase, V. V. Nesterenko and M. Bordag, J. Math. Phys. (1999), 6254 [hep-th/9812059].
8. G. Esposito, A. Y. Kamenshchik and K. Kirsten, Phys. Rev. D **62**, 085027-1 (2000) [hep-th/0006220].
9. G. Barton, J. Phys. A **37**, 1011 (2004).
10. E. M. Lifshitz, Zh. Eksp. Teor. Fiz. **29**, 94 (1956);
11. P. W. Millonni, The Quantum Vacuum (Academic, New York, 1993).

12. K. A. Milton, J. Phys. A **37**, R209 (2004) [hep-th/0406024].
13. R. B. Balian and C. Bloch, Ann. Phys. (NY) **60**, 401 (1970); *ibid* **63**, 592 (1971); *ibid* **64**, 271 (1971) Errata in *ibid* **84**, 559 (1974); *ibid* **69**, 76 (1972); *ibid* **85**, 514 (1974).
14. W. Lukosz, Physica **56**, 109 (1971).
15. J.S. Dowker and R. Critchley, J. Phys. A **9**, 535 (1976).
16. J. Ambjörn and S. Wolfram, Annals Phys. **147**, 1 (1983).
17. A. Edery, J. Phys. A **39**, 685 (2006).
18. I. L. DeRaad, Jr. and K. A. Milton, Ann. Phys. (N.Y.) **136**, 229 (1981).
19. P. Gosdzinsky and A. Romeo, Phys. Lett. B **476**, 369 (1998) [hep-th/9809199].
20. I. Brevik and G.H. Nyland, Ann. Phys. **230**, 321 (1994).
21. K.A. Milton, A.V. Nesterenko and V.V. Nesterenko, Phys. Rev. D **59**, 105009 (1999).
22. I. Klich and A. Romeo, Phys. Lett. B **476**, 369 (2000).
23. V. V. Nesterenko and I. G. Pirozhenko, J. Math. Phys. **41**, 4521 (2000) [hep-th/9910097].
24. R. Balian and B. Duplantier, Recent Developments in Gravitational Physics, Institute of Physics Conference Series **176**, Ed. Ciufolini et al, Oct. 2004 [quant-ph/0408124].
25. M. Schaden and L. Spruch, Phys. Rev. A **58**, 935 (1998).
26. M. C. Gutzwiller, *Chaos in Classical Quantum Mechanics* (Springer, Berlin, 1990) with references to the author's earlier work.
27. O. Schröder, A. Scardicchio, R. L. Jaffe, Phys. Rev. A **72**, 012105 (2005).
28. A. Bulgac, P. Magierski and A. Wirzba, Phys. Rev. D **73**, 025007 (2006).
29. M.V. Berry and M. Tabor, Proc.Roy.Soc.London **A349**, 101 (1976); *ibid* J. Phys. A **10**, 371 (1977); Proc.Roy.Soc.London **A356**, 375 (1977).
30. F. Mazzitelli, M.J. Sanchez, N.N. Scoccola, J. von Stecher, Phys. Rev. A **67**, 013807 (2003).
31. M. Schaden, Phys. Rev. A **73**, 042102 (2006).
32. C. Schubert, Phys. Rept. **355**, 73 (2002) [hep-th/0101036];
33. H. Gies and K. Langfeld, Int.J.Mod. Phys. A **17**, 966 (2002) [hep-th/0112198]; H. Gies, K. Langfeld and L. Moyaerts, JHEP **0306**, 018 (2003) [hep-th/0303264]; *ibid*, [hep-th/0311168]; H. Gies and K. Klingmuller, preprint quant-ph/0601094.
34. H. Gies and K. Klingmuller, Proceedings of QFEXT'05 (Barcelona, Sept. '05) hep-th/0511092; *ibid*, preprint quant-ph/0601094.
35. T. Emig, A. Hanke, R. Golestanian and M. Kardar, Phys. Rev. Lett. **87**, 260402 (2001); R. Büscher and T. Emig, Phys. Rev. Lett. **94**, 133901 (2005); T. Emig, R. L. Jaffe, M. Kardar and A. Scardicchio, preprint cond-mat/0601055.
36. G. Barton, J. Phys. A **34**, 4083 (2001).
37. I. Cavero-Peláez and K. A. Milton, Ann. Phys. (N.Y.) **320**, 108 (2005) [hep-th/0412135].
38. A. Romeo and K.A. Milton, Phys. Lett. B **621**, 309 (2005) [hep-th/0504207].
39. A. Einstein, Verh. Dtsch. Phys. Ges. **19**, 82 (1917).
40. J. B. Keller, Ann. Phys. (NY) **4**, 180 (1958); *ibid* in *Calculus of Variations and its Applications* (Am. Math. Soc., Providence, 1958), p.27; *ibid*, J. Opt. Soc. Am. **52**, 116 (1962).
41. V. P. Maslov, *Théorie des Perturbations et Méthodes Asymptotiques* (Dunod, Paris, 1972) and V. P. Maslov and M. V. Fedoriuk, *Semi-Classical Approximations in Quantum Mechanics* (Reidel, Boston, 1981).
42. D. Deutsch and P. Candelas, Phys. Rev. D **20**, 3063 (1979); P. Candelas, Ann. Phys. (NY) **143**, 241 (1982); *ibid*, **167**, 257 (1986).
43. J.S. Dowker, Ann. Phys. (N.Y.) **62**, 361 (1971).
44. P. Chang and J.S. Dowker, Nucl. Phys. B **395**, 407 (1993).
45. J.S. Dowker, preprint hep-th/0510248; *ibid*, preprint math.sp/0601334.
46. C. M. Bender, K. A. Milton, Phys. Rev. D **50**, 6547 (1994); K. A. Milton, Phys. Rev. D **55**, 4940 (1997).
47. A. A. Saharian, Preprint IC/2000/14, hep-th/0002239.
48. F. Bernasconi, G.M. Graf and D. Hasler,

- Ann. Inst. H. Poincaré **4**, 1001 (2003) [math-ph/0302035].
49. M. Bordag, K. Kirsten, D. Vassilevich, Phys. Rev. D **59**, 085011 (1999).
 50. N. Graham, R. L. Jaffe, V. Klemani, M. Quandt, M. Scandurra and H. Weigel, Nucl. Phys. B **645**, 49 (2002); R. L. Jaffe, AIP Conf. Proc. **687**, 3 (2003) [arXiv:hep-th/0307014].
 51. K. A. Milton, Phys. Rev. D **68**, 065020 (2003); *ibid* J. Phys. A **37**, R209 (2004).
 52. A.A. Saharian and A. S. Taroyan, hep-th/0603144.
 53. A. Scardicchio, R. L. Jaffe, Nucl. Phys. **704**, 552 (2005); *ibid*, preprint quant-ph/0507042.
 54. J.B. Keller, J. Opt. Soc. Amer. **52**, 116 (1962); J.B. Keller and R.M. Lewis, *Asymptotic Theory of Wave Propagation and Diffraction*, unpublished book manuscript in 2 Volumes, (Courant Institute, New York University, 1970); J.B. Keller, J. Appl. Phys. **30**, 1452 (1952); B.R. Levy and J. Keller, Commun. Pure Appl. Math. **12**, 159 (1959); H.M. Nussenzveig, Ann. Phys. **34**, 23 (1965); H. Primack, H. Schranz, U. Smilansky and I. Ussishkin, J. Phys. A: Math. Gen. **30**, 6693 (1997).
 55. M. Schaden and L. Spruch, Ann. Phys. (N.Y.) **313**, 37 (2004).
 56. G. Vattay, A. Wirzba, P.E. Rosenqvist, Phys. Rev. Lett. **73** (1994) 2304.
 57. A. Romeo, private communication to K. A. Milton (reproduced in App. B of [21]).
 58. M. Bordag and I. G. Pirozhenko, Phys. Rev. D **64**, 025019 (2001).
 59. V.V. Nesterenko and I. G. Pirozhenko, Phys. Rev. D **60**, 125007 (1999).
 60. E. A. Power, *Introductory Quantum Electrodynamics* (Elsevier, New York, 1964), Appendix I.
 61. L. Brown and G. J. Maclay, Phys. Rev. **184**, 1272 (1969); M. Schaden and L. Spruch, Phys. Rev. A **65**, 034101 (2002).
 62. J. S. Høye, I. Brevik, J. B. Aarseth, K. A. Milton, quant-ph/0506025.
 63. P.M. Neto, A. Lambrecht and S. Reynaud, quant-ph/0505086.